

Two dimensional event set and its application in algebraic structures

Y.B. Jun¹

¹Department of Mathematics Education, Gyeongsang National University, Jinju 52828, Korea

skywine@gmail.com

Abstract

Two dimensional event set is introduced, and it is applied to algebraic structures. Two dimensional BCK/BCI-eventful algebra, paired B-algebra and paired BCK/BCI-algebra are defined, and several properties are investigated. Conditions for two dimensional eventful algebra to be a B-algebra and a BCK/BCI-algebra are provided. The process of inducing a paired B-algebra using a group is discussed. Using two dimensional BCI-eventful algebra, a commutative group is established.

Article Information

Corresponding Author:

Y.B. Jun;

Received: March 2020 ;

Accepted: Invited paper.

Keywords:

Two dimensional event (set), two dimensional BCK/BCI-eventful algebra, paired B-algebra, paired BCK/BCI-algebra.



Check for updates

1 Introduction and Preliminaries

The notion of neutrosophic set is developed by Smarandache ([9], [10]), and is a more general platform that extends the notions of classic set, (intuitionistic) fuzzy set and interval valued (intuitionistic) fuzzy set. Smarandache [11] considered an entry (i.e., a number, an idea, an object etc.) which is represented by a known part (a) and an unknown part (bT, cI, dF) where T, I, F have their usual neutrosophic logic meanings and a, b, c, d are real or complex numbers, and then he introduced the concept of neutrosophic quadruple numbers. Neutrosophic quadruple algebraic structures and hyperstructures are discussed in [1] and [2]. Neutrosophic quadruple BCK/BCI-algebra is studied in [5] and [7]. Using neutrosophic quadruple structures, Jun et al. [4] introduced the notion of events by considering two facts, and applied it to BCK/BCI-algebras. There are many things in our daily lives that we have to choose between two facts. For example, should I read a book or not, go to the movies or not, etc. To consider these two factors, we introduce two-dimensional event sets and try to apply them to algebraic structures. We introduce the notions of two dimensional BCK/BCI-eventful algebra, paired B-algebra and paired BCK/BCI-algebra, and investigate several properties. We provided conditions for two dimensional eventful algebra to be a B-algebra and a BCK/BCI-algebra. We discuss the process of inducing a paired B-algebra using a group, and establish a commutative group using two dimensional BCI-eventful algebra.

We describe the basic contents that will be needed in this paper. Let $(X, *, 0)$ be an algebra, i.e., let X be a set with a special element “0” and a binary operation “*”, and consider the following conditions.

$$(\forall u \in X) (u * u = 0). \quad (1)$$

$$(\forall u \in X) (u * 0 = u). \quad (2)$$

$$(\forall u \in X) (0 * u = 0). \quad (3)$$

$$(\forall u, v, w \in X) ((u * v) * w = u * (w * (0 * v))). \quad (4)$$

$$(\forall u, v \in X) (u * v = 0, v * u = 0 \Rightarrow u = v). \quad (5)$$

$$(\forall u, v \in X) ((u * (u * v)) * v = 0). \quad (6)$$

$$(\forall u, v, w \in X) (((u * v) * (u * w)) * (w * v) = 0). \quad (7)$$

We say that $X := (X, *, 0)$ is

- a *B-algebra* (see [8]) if it satisfies (1), (2) and (4),
- a *BCI-algebra* (see [6]) if it satisfies (1), (5), (6) and (7),
- a *BCK-algebra* (see [3]) if it is a BCI-algebra satisfying (3).

A BCI-algebra $X := (X, *, 0)$ is said to be *p-semisimple* (see [3]) if it satisfies:

$$(\forall u \in X) (0 * (0 * u) = u). \quad (8)$$

2 Two dimensional event sets

Definition 2.1. Let $\ell : X \rightarrow Q$ be a mapping from a set X to a set Q . For any $a, x \in X$, the ordered pair (x, ℓ_a) is called a two dimensional event on X where ℓ_a is the image of a under ℓ .

The set of all two dimensional events on X is denoted by (X, ℓ_X) , that is,

$$(X, \ell_X) = \{(x, \ell_a) \mid x, a \in X\} \quad (9)$$

and it is called a *two dimensional X-event set*. By a *two dimensional X-eventful algebra* we mean a two dimensional X -event set with a binary operation $\&$, and it is denoted by $\langle (X, \ell_X), \& \rangle$.

Let $\langle (\mathbb{R}, \ell_{\mathbb{R}}), \oplus \rangle$, $\langle (\mathbb{R}, \ell_{\mathbb{R}}), \ominus \rangle$ and $\langle (\mathbb{R}, \ell_{\mathbb{R}}), \odot \rangle$ be two dimensional \mathbb{R} -eventful algebras in which “ \oplus ”, “ \ominus ” and “ \odot ” are defined as follows:

$$(x, \ell_a) \oplus (y, \ell_b) = (x + y, \ell_{a+b}),$$

$$(x, \ell_a) \ominus (y, \ell_b) = (x - y, \ell_{a-b}),$$

$$(x, \ell_a) \odot (y, \ell_b) = (x \cdot y, \ell_{a \cdot b}),$$

respectively, for any two dimensional events (x, ℓ_a) and (y, ℓ_b) on \mathbb{R} . For any $t \in \mathbb{R}$ and a two dimensional event (x, ℓ_a) on \mathbb{R} , we define

$$t(x, \ell_a) = (tx, \ell_{ta}). \quad (10)$$

In particular, if $t = -1$, then $-1(x, \ell_a) = (-x, \ell_{-a})$ and $-1(x, \ell_a)$ is simply denoted by $-(x, \ell_a)$.

Proposition 2.2. Let $\langle (\mathbb{R}, \ell_{\mathbb{R}}), \oplus \rangle$, $\langle (\mathbb{R}, \ell_{\mathbb{R}}), \ominus \rangle$ and $\langle (\mathbb{R}, \ell_{\mathbb{R}}), \odot \rangle$ be two dimensional \mathbb{R} -eventful algebras. Then

$$(i) \ ((x, \ell_a) \oplus (y, \ell_b)) \oplus (z, \ell_c) = (x, \ell_a) \oplus ((y, \ell_b) \oplus (z, \ell_c)),$$

$$(ii) \ (x, \ell_a) \oplus (y, \ell_b) = (y, \ell_b) \oplus (x, \ell_a),$$

$$(iii) \ (x, \ell_a) \odot (y, \ell_b) = (y, \ell_b) \odot (x, \ell_a),$$

$$(iv) ((x, \ell_a) \odot (y, \ell_b)) \odot (z, \ell_c) = (x, \ell_a) \odot ((y, \ell_b) \odot (z, \ell_c)),$$

$$(v) t((x, \ell_a) \oplus (y, \ell_b)) = t(x, \ell_a) \oplus t(y, \ell_b) \text{ for all } t \in \mathbb{R},$$

$$(vi) (t + s)(x, \ell_a) = t(x, \ell_a) \oplus s(x, \ell_a) \text{ for all } t, s \in \mathbb{R},$$

$$(vii) (x, \ell_a) \oplus (-(x, \ell_a)) = (0, \ell_0).$$

$$(viii) (x, \ell_a) \odot (x, \ell_a)^{-1} = (1, \ell_1) \text{ where } (x, \ell_a)^{-1} = (x^{-1}, \ell_{a^{-1}})$$

for all two dimensional events (x, ℓ_a) , (y, ℓ_b) and (z, ℓ_c) on \mathbb{R} .

Proof. Straightforward. □

By Proposition 2.2, we have the following theorem.

Theorem 2.3. *Two dimensional \mathbb{R} -eventful algebras $\langle(\mathbb{R}, \ell_{\mathbb{R}}), \oplus\rangle$ and $\langle(\mathbb{R}, \ell_{\mathbb{R}}), \odot\rangle$ are commutative groups with identities $(0, \ell_0)$ and $(1, \ell_1)$, respectively.*

3 Two dimensional eventful algebras

Let $(X, *, 0)$ be an algebra and $\langle(X, \ell_X), \otimes\rangle$ be a two dimensional X -eventful algebra in which “ \otimes ” is defined by

$$(x, \ell_a) \otimes (y, \ell_b) = (x * y, \ell_{a*b})$$

respectively, for all $x, y, a, b \in X$. In a two dimensional X -eventful algebra $\langle(X, \ell_X), \otimes\rangle$, the order “ \ll ” is defined as follows:

$$(x, \ell_a) \ll (y, \ell_b) \Leftrightarrow x \leq y \text{ and } a \leq b$$

for all $x, y, a, b \in X$ where $x \leq y$ means $x * y = 0$ and $a \leq b$ means $a * b = 0$.

Theorem 3.1. *If $(X, *, 0)$ is a B-algebra, then the two dimensional X -eventful algebra $\langle(X, \ell_X), \otimes\rangle$ is a B-algebra with the special element $(0, \ell_0)$.*

Proof. For any $(x, \ell_a), (y, \ell_b), (z, \ell_c) \in (X, \ell_X)$, we have

$$(x, \ell_a) \otimes (x, \ell_a) = (x * x, \ell_{a*a}) = (0, \ell_0), (x, \ell_a) \otimes (0, \ell_0) = (x * 0, \ell_{a*0}) = (x, \ell_a),$$

and

$$\begin{aligned} ((x, \ell_a) \otimes (y, \ell_b)) \otimes (z, \ell_c) &= (x * y, \ell_{a*b}) \otimes (z, \ell_c) \\ &= ((x * y) * z, \ell_{(a*b)*c}) \\ &= (x * (z * (0 * y)), \ell_{a*(c*(0*b))}) \\ &= (x, \ell_a) \otimes (z * (0 * y), \ell_{c*(0*b)}) \\ &= (x, \ell_a) \otimes ((z, \ell_c) \otimes (0 * y, \ell_{0*b})) \\ &= (x, \ell_a) \otimes ((z, \ell_c) \otimes ((0, \ell_0) \otimes (y, \ell_b))) \end{aligned}$$

by (1), (2) and (4), respectively. Therefore $\langle(X, \ell_X), \otimes\rangle$ is a B-algebra with the special element $(0, \ell_0)$. □

We say that $\langle(X, \ell_X), \otimes, (0, \ell_0)\rangle$ is a *paired B-algebra*.

Example 3.2. *Let $X = \{0, a, b\}$ be a set with the binary operation “ $*$ ” as in Table 1.*

For $Q = \{0, \frac{1}{2}, 1\}$, define a mapping ℓ as follows:

$$\ell : X \rightarrow Q, x \mapsto \begin{cases} 0 & \text{if } x = 0, \\ \frac{1}{2} & \text{if } x = a, \\ 1 & \text{if } x = b. \end{cases}$$

Table 1: Cayley table for the binary operation “*”

*	0	a	b
0	0	b	a
a	a	0	b
b	b	a	0

Then

$$\begin{aligned} (X, \ell_X) &= \{(0, \ell_0), (0, \ell_a), (0, \ell_b), (a, \ell_0), (a, \ell_a), (a, \ell_b), (b, \ell_0), (b, \ell_a), (b, \ell_b)\} \\ &= \{(0, 0), (0, \frac{1}{2}), (0, 1), (a, 0), (a, \frac{1}{2}), (a, 1), (b, 0), (b, \frac{1}{2}), (b, 1)\} \end{aligned}$$

and the operation \otimes is given by Table 2. It is routine to verify that $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ is a paired B -algebra.

Table 2: Cayley table for the binary operation “*”

\otimes	$(0, \ell_0)$	$(0, \ell_a)$	$(0, \ell_b)$	(a, ℓ_0)	(a, ℓ_a)	(a, ℓ_b)	(b, ℓ_0)	(b, ℓ_a)	(b, ℓ_b)
$(0, \ell_0)$	$(0, 0)$	$(0, 1)$	$(0, \frac{1}{2})$	$(b, 0)$	$(b, 1)$	$(b, \frac{1}{2})$	$(a, 0)$	$(a, 1)$	$(a, \frac{1}{2})$
$(0, \ell_a)$	$(0, \frac{1}{2})$	$(0, 0)$	$(0, 1)$	$(b, \frac{1}{2})$	$(b, 0)$	$(b, 1)$	$(a, \frac{1}{2})$	$(a, 0)$	$(a, 1)$
$(0, \ell_b)$	$(0, 1)$	$(0, \frac{1}{2})$	$(0, 0)$	$(b, 1)$	$(b, \frac{1}{2})$	$(b, 0)$	$(a, 1)$	$(a, \frac{1}{2})$	$(a, 0)$
(a, ℓ_0)	$(a, 0)$	$(a, 1)$	$(a, \frac{1}{2})$	$(0, 0)$	$(0, 1)$	$(0, \frac{1}{2})$	$(b, 0)$	$(b, 1)$	$(b, \frac{1}{2})$
(a, ℓ_a)	$(a, \frac{1}{2})$	$(a, 0)$	$(a, 1)$	$(0, \frac{1}{2})$	$(0, 0)$	$(0, 1)$	$(b, \frac{1}{2})$	$(b, 0)$	$(b, 1)$
(a, ℓ_b)	$(a, 1)$	$(a, \frac{1}{2})$	$(a, 0)$	$(0, 1)$	$(0, \frac{1}{2})$	$(0, 0)$	$(b, 1)$	$(b, \frac{1}{2})$	$(b, 0)$
(b, ℓ_0)	$(b, 0)$	$(b, 1)$	$(b, \frac{1}{2})$	$(a, 0)$	$(a, 1)$	$(a, \frac{1}{2})$	$(0, 0)$	$(0, 1)$	$(0, \frac{1}{2})$
(b, ℓ_a)	$(b, \frac{1}{2})$	$(b, 0)$	$(b, 1)$	$(a, \frac{1}{2})$	$(a, 0)$	$(a, 1)$	$(0, \frac{1}{2})$	$(0, 0)$	$(0, 1)$
(b, ℓ_b)	$(b, 1)$	$(b, \frac{1}{2})$	$(b, 0)$	$(a, 1)$	$(a, \frac{1}{2})$	$(a, 0)$	$(0, 1)$	$(0, \frac{1}{2})$	$(0, 0)$

Proposition 3.3. *If $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ is a paired B -algebra, then*

- (i) $(x, \ell_a) \otimes (y, \ell_b) = (x, \ell_a) \otimes ((0, \ell_0) \otimes ((0, \ell_0) \otimes (y, \ell_b)))$,
- (ii) $((x, \ell_a) \otimes (y, \ell_b)) \otimes ((0, \ell_0) \otimes (y, \ell_b)) = (x, \ell_a)$,
- (iii) $(x, \ell_a) \otimes (z, \ell_c) = (y, \ell_b) \otimes (z, \ell_c)$ implies $(x, \ell_a) = (y, \ell_b)$,
- (iv) $(x, \ell_a) \otimes ((y, \ell_b) \otimes (z, \ell_c)) = ((x, \ell_a) \otimes ((0, \ell_0) \otimes (z, \ell_c))) \otimes (y, \ell_b)$,
- (v) $(x, \ell_a) \otimes (y, \ell_b) = (0, \ell_0)$ implies $(x, \ell_a) = (y, \ell_b)$,
- (vi) $(0, \ell_0) \otimes (x, \ell_a) = (0, \ell_0) \otimes (y, \ell_b)$ implies $(x, \ell_a) = (y, \ell_b)$,
- (vii) $(0, \ell_0) \otimes ((0, \ell_0) \otimes (x, \ell_a)) = (x, \ell_a)$

for all $x, y, z, a, b, c \in X$.

Proof. Let $x, y, z, a, b, c \in X$. Then

$$\begin{aligned} (x, \ell_a) \otimes (y, \ell_b) &= (x * y, \ell_{a*b}) = ((x * y) * 0, \ell_{(a*b)*0}) \\ &= (x * (0 * (0 * y)), \ell_{a*(0*(0*b))}) \\ &= (x, \ell_a) \otimes (0 * (0 * y), \ell_{0*(0*b)}) \\ &= (x, \ell_a) \otimes ((0, \ell_0) \otimes ((0, \ell_0) \otimes (y, \ell_b))) \end{aligned}$$

which proves (i).

(ii) We have

$$\begin{aligned}
& ((x, \ell_a) \otimes (y, \ell_b)) \otimes ((0, \ell_0) \otimes (y, \ell_b)) = (x * y, \ell_{a*b}) \otimes (0 * y, \ell_{0*b}) \\
& = ((x * y) * (0 * y), \ell_{(a*b)*(0*b)}) \\
& = (x * ((0 * y) * (0 * y)), \ell_{a*((0*b)*(0*b))}) \\
& = (x * 0, \ell_{a*0}) = (x, \ell_a).
\end{aligned}$$

(iii) Assume that $(x, \ell_a) \otimes (z, \ell_c) = (y, \ell_b) \otimes (z, \ell_c)$. It follows from (ii) that

$$\begin{aligned}
(x, \ell_a) &= ((x, \ell_a) \otimes (z, \ell_c)) \otimes ((0, \ell_0) \otimes (z, \ell_c)) \\
&= ((y, \ell_b) \otimes (z, \ell_c)) \otimes ((0, \ell_0) \otimes (z, \ell_c)) \\
&= (y, \ell_b).
\end{aligned}$$

(iv) Using (i), we have

$$\begin{aligned}
& ((x, \ell_a) \otimes ((0, \ell_0) \otimes (z, \ell_c))) \otimes (y, \ell_b) \\
&= (x, \ell_a) \otimes ((y, \ell_b) \otimes ((0, \ell_0) \otimes ((0, \ell_0) \otimes (z, \ell_c)))) \\
&= (x, \ell_a) \otimes ((y, \ell_b) \otimes (z, \ell_c)).
\end{aligned}$$

(v) Suppose that $(x, \ell_a) \otimes (y, \ell_b) = (0, \ell_0)$. Then $(x, \ell_a) \otimes (y, \ell_b) = (y, \ell_b) \otimes (y, \ell_b)$, and so $(x, \ell_a) = (y, \ell_b)$ by (iii).

(vi) Assume that $(0, \ell_0) \otimes (x, \ell_a) = (0, \ell_0) \otimes (y, \ell_b)$. Then

$$\begin{aligned}
(0, \ell_0) &= (x, \ell_a) \otimes (x, \ell_a) = (x, \ell_a) \otimes ((0, \ell_0) \otimes ((0, \ell_0) \otimes (x, \ell_a))) \\
&= (x, \ell_a) \otimes ((0, \ell_0) \otimes ((0, \ell_0) \otimes (y, \ell_b))) \\
&= (x, \ell_a) \otimes (y, \ell_b)
\end{aligned}$$

and so $(x, \ell_a) = (y, \ell_b)$ by (v).

(vii) We have

$$\begin{aligned}
(0, \ell_0) \otimes (x, \ell_a) &= ((0, \ell_0) \otimes (x, \ell_a)) \otimes (0, \ell_0) \\
&= ((0 * x) * 0, \ell_{(0*a)*0}) \\
&= (0 * (0 * (0 * x)), \ell_{0*(0*(0*a))}) \\
&= (0, \ell_0) \otimes ((0, \ell_0) \otimes ((0, \ell_0) \otimes (x, \ell_a)))
\end{aligned}$$

and so $(0, \ell_0) \otimes ((0, \ell_0) \otimes (x, \ell_a)) = (x, \ell_a)$ by (vi). □

We provide conditions for two dimensional X -eventful algebra to be a B-algebra.

Theorem 3.4. *For an algebra $(X, *, 0)$, the two dimensional X -eventful algebra $\langle (X, \ell_X), \otimes \rangle$ is a B-algebra with the special element $(0, \ell_0)$ if and only if it satisfies Proposition 3.3(vii) and*

$$(x, \ell_a) \otimes (x, \ell_a) = (0, \ell_0), \tag{11}$$

$$((x, \ell_a) \otimes (z, \ell_c)) \otimes ((y, \ell_b) \otimes (z, \ell_c)) = (x, \ell_a) \otimes (y, \ell_b) \tag{12}$$

for all $(x, \ell_a), (y, \ell_b), (z, \ell_c) \in (X, \ell_X)$.

Proof. Assume that the two dimensional X -eventful algebra $\langle (X, \ell_X), \otimes \rangle$ is a B-algebra with the special element $(0, \ell_0)$. The condition Proposition 3.3(vii) is by Proposition 3.3. It is clear that (11) is true by the definition of B-algebra. Also, we have

$$\begin{aligned}
& ((x, \ell_a) \otimes (z, \ell_c)) \otimes ((y, \ell_b) \otimes (z, \ell_c)) = (x, \ell_a) \otimes (((y, \ell_b) \otimes (z, \ell_c)) \otimes ((0, \ell_0) \otimes (z, \ell_c))) \\
&= (x, \ell_a) \otimes ((y, \ell_b) \otimes (((0, \ell_0) \otimes (z, \ell_c)) \otimes ((0, \ell_0) \otimes (z, \ell_c)))) \\
&= (x, \ell_a) \otimes ((y, \ell_b) \otimes (0, \ell_0)) = (x, \ell_a) \otimes (y, \ell_b)
\end{aligned}$$

for all $(x, \ell_a), (y, \ell_b), (z, \ell_c) \in (X, \ell_X)$.

Conversely, suppose that $\langle (X, \ell_X), \otimes \rangle$ satisfies three conditions (11), (12) and Proposition 3.3(vii). Then

$$\begin{aligned} (x, \ell_a) &= (0, \ell_0) \otimes ((0, \ell_0) \otimes (x, \ell_a)) \\ &= ((x, \ell_a) \otimes (x, \ell_a)) \otimes ((0, \ell_0) \otimes (x, \ell_a)) \\ &= (x, \ell_a) \otimes (0, \ell_0) \end{aligned}$$

and

$$((x, \ell_a) \otimes (y, \ell_b)) \otimes ((0, \ell_0) \otimes (y, \ell_b)) = (x, \ell_a). \quad (13)$$

Combining (12) with (13) induces

$$\begin{aligned} &(x, \ell_a) \otimes ((z, \ell_c) \otimes ((0, \ell_0) \otimes (y, \ell_b))) \\ &= (((x, \ell_a) \otimes (y, \ell_b)) * ((0, \ell_0) \otimes (y, \ell_b))) \otimes ((z, \ell_c) \otimes ((0, \ell_0) \otimes (y, \ell_b))) \\ &= ((x, \ell_a) \otimes (y, \ell_b)) \otimes (z, \ell_c) \end{aligned}$$

for all $(x, \ell_a), (y, \ell_b), (z, \ell_c) \in (X, \ell_X)$. Therefore $\langle (X, \ell_X), \otimes \rangle$ is a B-algebra with the special element $(0, \ell_0)$. \square

Theorem 3.5. *For an algebra $(X, *, 0)$, the two dimensional X -eventful algebra $\langle (X, \ell_X), \otimes \rangle$ is a B-algebra with the special element $(0, \ell_0)$ if and only if it satisfies (11) and*

$$(x, \ell_a) \otimes (((0, \ell_0) \otimes (y, \ell_b)) \otimes (z, \ell_c)) \otimes (((0, \ell_0) \otimes (x, \ell_a)) \otimes (z, \ell_c)) = (y, \ell_b) \quad (14)$$

for all $(x, \ell_a), (y, \ell_b), (z, \ell_c) \in (X, \ell_X)$.

Proof. Assume that the two dimensional X -eventful algebra $\langle (X, \ell_X), \otimes \rangle$ is a B-algebra with the special element $(0, \ell_0)$. Then (11) is valid in Theorem 3.4. Using (12), we get

$$\begin{aligned} &(((0, \ell_0) \otimes (y, \ell_b)) \otimes (z, \ell_c)) \otimes (((0, \ell_0) \otimes (x, \ell_a)) \otimes (z, \ell_c)) \\ &= ((0, \ell_0) \otimes (y, \ell_b)) \otimes ((0, \ell_0) \otimes (x, \ell_a)). \end{aligned} \quad (15)$$

It follows that

$$\begin{aligned} &(x, \ell_a) \otimes (((0, \ell_0) \otimes (y, \ell_b)) \otimes (z, \ell_c)) \otimes (((0, \ell_0) \otimes (x, \ell_a)) \otimes (z, \ell_c)) \\ &= (x, \ell_a) \otimes (((0, \ell_0) \otimes (y, \ell_b)) \otimes ((0, \ell_0) \otimes (x, \ell_a))) \\ &= ((x, \ell_a) \otimes (x, \ell_a)) \otimes ((0, \ell_0) \otimes (y, \ell_b)) \\ &= (0, \ell_0) \otimes ((0, \ell_0) \otimes (y, \ell_b)) \\ &= (y, \ell_b) \end{aligned}$$

which proves (14).

Conversely, suppose that $\langle (X, \ell_X), \otimes \rangle$ satisfies (11) and (14). If we substitute (y, ℓ_b) for (x, ℓ_a) in (14) and use (11), then

$$\begin{aligned} (x, \ell_a) &= (x, \ell_a) \otimes (((0, \ell_0) \otimes (x, \ell_a)) \otimes (z, \ell_c)) \otimes (((0, \ell_0) \otimes (x, \ell_a)) \otimes (z, \ell_c)) \\ &= (x, \ell_a) \otimes (0, \ell_0) \end{aligned} \quad (16)$$

If we put $(x, \ell_a) = (0, \ell_0) = (z, \ell_c)$ and $(y, \ell_b) = (x, \ell_a)$ in (14), then

$$\begin{aligned} (x, \ell_a) &= (0, \ell_0) \otimes (((0, \ell_0) \otimes (x, \ell_a)) \otimes (0, \ell_0)) \otimes (((0, \ell_0) \otimes (0, \ell_0)) \otimes (0, \ell_0)) \\ &= (0, \ell_0) \otimes ((0, \ell_0) \otimes (x, \ell_a)) \end{aligned} \quad (17)$$

by (16). Assume that $(0, \ell_0) \otimes (x, \ell_a) = (0, \ell_0) \otimes (y, \ell_b)$. Then

$$(x, \ell_a) = (0, \ell_0) \otimes ((0, \ell_0) \otimes (x, \ell_a)) = (0, \ell_0) \otimes ((0, \ell_0) \otimes (y, \ell_b)) = (y, \ell_b)$$

by (17) which proves

$$(0, \ell_0) \otimes (x, \ell_a) = (0, \ell_0) \otimes (y, \ell_b) \Rightarrow (x, \ell_a) = (y, \ell_b). \quad (18)$$

Putting $(x, \ell_a) = (0, \ell_0)$, $(y, \ell_b) = (0, \ell_0) \otimes (y', \ell_{b'})$ and $(z, \ell_c) = (z', \ell_{c'})$ in (14) induces

$$\begin{aligned} (0, \ell_0) \otimes (y', \ell_{b'}) &= (0, \ell_0) \otimes (((0, \ell_0) \otimes ((0, \ell_0) \otimes (y', \ell_{b'})) \otimes (z', \ell_{c'})) \otimes \\ &\quad (((0, \ell_0) \otimes (0, \ell_0)) \otimes (z', \ell_{c'}))) \\ &= (0, \ell_0) \otimes ((y', \ell_{b'}) \otimes (z', \ell_{c'})) \otimes ((0, \ell_0) \otimes (z', \ell_{c'})). \end{aligned}$$

It follows from (18) that

$$(y', \ell_{b'}) = (((y', \ell_{b'}) \otimes (z', \ell_{c'})) \otimes ((0, \ell_0) \otimes (z', \ell_{c'}))). \quad (19)$$

If we substitute (x, ℓ_a) , (y, ℓ_b) and (z, ℓ_c) for $(x', \ell_{a'})$, $(0, \ell_0) \otimes ((z', \ell_{c'}) \otimes (x', \ell_{a'}))$ and $(0, \ell_0)$, respectively, in (14), then

$$\begin{aligned} (0, \ell_0) \otimes ((z', \ell_{c'}) \otimes (x', \ell_{a'})) &= (x', \ell_{a'}) \otimes (((0, \ell_0) \otimes ((0, \ell_0) \otimes ((z', \ell_{c'}) \otimes (x', \ell_{a'})))) \otimes (0, \ell_0) \\ &\quad \otimes (((0, \ell_0) \otimes (x', \ell_{a'})) \otimes (0, \ell_0)) \\ &= (x', \ell_{a'}) \otimes (((z', \ell_{c'}) \otimes (x', \ell_{a'})) \otimes ((0, \ell_0) \otimes (x', \ell_{a'}))) \\ &= ((x', \ell_{a'}) \otimes (z', \ell_{c'})) \end{aligned} \quad (20)$$

In (14), taking $(x, \ell_a) = (y', \ell_{b'}) \otimes (w, \ell_d)$, $(y, \ell_b) = (y', \ell_{b'}) \otimes (z', \ell_{c'})$ and $(z, \ell_c) = (0, \ell_0) \otimes (y', \ell_{b'})$ imply that

$$\begin{aligned} (y', \ell_{b'}) \otimes (z', \ell_{c'}) &= ((y', \ell_{b'}) \otimes (w, \ell_d)) \otimes [(((0, \ell_0) \otimes ((y', \ell_{b'}) \otimes (z', \ell_{c'}))) \\ &\quad \otimes ((0, \ell_0) \otimes (y', \ell_{b'}))) \otimes (((0, \ell_0) \otimes ((y', \ell_{b'}) \otimes (w, \ell_d))) \\ &\quad \otimes ((0, \ell_0) \otimes (y', \ell_{b'})))] \end{aligned} \quad (21)$$

Using (19) and (20), we get

$$\begin{aligned} ((0, \ell_0) \otimes ((y', \ell_{b'}) \otimes (z', \ell_{c'}))) \otimes ((0, \ell_0) \otimes (y', \ell_{b'})) &= ((z', \ell_{c'}) \otimes (y', \ell_{b'})) \otimes ((0, \ell_0) \otimes (y', \ell_{b'})) = (z', \ell_{c'}). \end{aligned} \quad (22)$$

Similarly, we have

$$((0, \ell_0) \otimes ((y', \ell_{b'}) \otimes (w, \ell_d))) \otimes ((0, \ell_0) \otimes (y', \ell_{b'})) = (w, \ell_d). \quad (23)$$

Combining (21), (22) and (23) induces

$$(y', \ell_{b'}) \otimes (z', \ell_{c'}) = ((y', \ell_{b'}) \otimes (w, \ell_d)) \otimes ((z', \ell_{c'}) \otimes (w, \ell_d)).$$

Therefore $\langle (X, \ell_X), \otimes \rangle$ is a B-algebra with the special element $(0, \ell_0)$ by Theorem 3.4. \square

The following theorem shows the process of inducing a paired B-algebra using a group.

Theorem 3.6. *If $(X, \circ, 0)$ is a group, then the two dimensional X-eventful algebra $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ is a paired B-algebra where*

$$\otimes : (X, \ell_X) \times (X, \ell_X) \rightarrow (X, \ell_X), ((x, \ell_a), (y, \ell_b)) \mapsto (x \circ y^{-1}, \ell_{a \circ b^{-1}}) = (x * y, \ell_{a * b}).$$

Proof. Let $(x, \ell_a), (y, \ell_b), (z, \ell_c) \in (X, \ell_X)$. Then $(x, \ell_a) \otimes (x, \ell_a) = (x \circ x^{-1}, \ell_{a \circ a^{-1}}) = (0, \ell_0)$ and $(x, \ell_a) \otimes (0, \ell_0) = (x \circ 0^{-1}, \ell_{a \circ 0^{-1}}) = (x \circ 0, \ell_{a \circ 0}) = (x, \ell_a)$. Also

$$\begin{aligned} ((x, \ell_a) \otimes (y, \ell_b)) \otimes (z, \ell_c) &= (x \circ y^{-1}, \ell_{a \circ b^{-1}}) \otimes (z, \ell_c) \\ &= ((x \circ y^{-1}) \circ z^{-1}, \ell_{(a \circ b^{-1}) \circ c^{-1}}) \\ &= (x \circ (z \circ y)^{-1}, \ell_{a \circ (c \circ b)^{-1}}) \end{aligned}$$

and

$$\begin{aligned}
(x, \ell_a) \otimes ((z, \ell_c) \otimes ((0, \ell_0) \otimes (y, \ell_b))) &= (x, \ell_a) \otimes ((z, \ell_c) \otimes (0 \circ y^{-1}, \ell_{0 \circ b^{-1}})) \\
&= (x, \ell_a) \otimes ((z, \ell_c) \otimes (y^{-1}, \ell_{b^{-1}})) = (x, \ell_a) \otimes (z \circ (y^{-1})^{-1}, \ell_{c \circ (b^{-1})^{-1}}) \\
&= (x, \ell_a) \otimes (z \circ y, \ell_{c \circ b}) = (x \circ (z \circ y)^{-1}, \ell_{a \circ (c \circ b)^{-1}}).
\end{aligned}$$

Hence $((x, \ell_a) \otimes (y, \ell_b)) \otimes (z, \ell_c) = (x, \ell_a) \otimes ((z, \ell_c) \otimes ((0, \ell_0) \otimes (y, \ell_b)))$. Therefore $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ is a paired B-algebra. \square

Let $X := (X, *, 0)$ be an algebra. In a two dimensional X -eventful algebra $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$, we consider the following assertions.

$$(((x, \ell_a) \otimes (y, \ell_b)) \otimes ((x, \ell_a) \otimes (z, \ell_c))) \otimes ((z, \ell_c) \otimes (y, \ell_b)) = (0, \ell_0), \quad (24)$$

$$((x, \ell_a) \otimes ((x, \ell_a) \otimes (y, \ell_b))) \otimes (y, \ell_b) = (0, \ell_0), \quad (25)$$

$$(0, \ell_0) \otimes (x, \ell_a) = (0, \ell_0), \quad (26)$$

$$(x, \ell_a) \otimes (y, \ell_b) = (0, \ell_0), (y, \ell_b) \otimes (x, \ell_a) = (0, \ell_0) \Rightarrow (x, \ell_a) = (y, \ell_b) \quad (27)$$

for all $x, y, z, a, b, c \in X$.

Definition 3.7. Given an algebra $X := (X, *, 0)$, let $\langle (X, \ell_X), \otimes \rangle$ be a two dimensional X -eventful algebra with a special element $(0, \ell_0)$. If it satisfies (11), (24) and (25), we say that $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ is a two dimensional BCI-eventful algebra. If a two dimensional BCI-eventful algebra $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ satisfies the condition (26), it is called a two dimensional BCK-eventful algebra.

Example 3.8. (1) Consider an algebra $X := (X, *, 0)$ where $X = \{0, a\}$ and the binary operation “*” is given by Table 3. Given a set $Q = \{\alpha, \beta\}$, define a mapping ℓ as follows:

Table 3: Cayley table for the binary operation “*”

*	0	a
0	0	0
a	a	0

$$\ell : X \rightarrow Q, x \mapsto \begin{cases} \alpha & \text{if } x = 0, \\ \beta & \text{if } x = a. \end{cases}$$

Then $(X, \ell_X) = \{(0, \ell_0), (0, \ell_a), (a, \ell_0), (a, \ell_a)\} = \{(0, \alpha), (0, \beta), (a, \alpha), (a, \beta)\}$ and the operation \otimes is given by Table 4. It is routine to verify that $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ is a two dimensional BCK-eventful algebra.

Table 4: Cayley table for the binary operation “*”

\otimes	$(0, \ell_0)$	$(0, \ell_a)$	(a, ℓ_0)	(a, ℓ_a)
$(0, \ell_0)$	$(0, \alpha)$	$(0, \alpha)$	$(0, \alpha)$	$(0, \alpha)$
$(0, \ell_a)$	$(0, \beta)$	$(0, \alpha)$	$(0, \beta)$	$(0, \alpha)$
(a, ℓ_0)	(a, α)	(a, α)	$(0, \alpha)$	$(0, \alpha)$
(a, ℓ_a)	(a, β)	(a, α)	$(0, \beta)$	$(0, \alpha)$

(2) The paired B-algebra $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ in Example 3.2 is two dimensional BCI-eventful algebra.

In general, two dimensional BCK/BCI-eventful algebra $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ does not satisfy the condition (27) as seen in the following examples.

Table 5: Cayley table for the binary operation “*”

*	0	a	b	c
0	0	0	c	b
a	a	0	c	b
b	b	b	0	c
c	c	c	b	0

Example 3.9. Consider an algebra $X = (X, *, 0)$ where $X = \{0, a, b, c\}$ and the binary operation “*” is given by Table 5.

Given a set $Q = \{0.2, 0.5, 0.7\}$, define a mapping ℓ as follows:

$$\ell : X \rightarrow Q, x \mapsto \begin{cases} 0.2 & \text{if } x \in \{0, a\} \\ 0.7 & \text{if } x \in \{b, c\}. \end{cases}$$

Then the two dimensional X -event set is given as follows:

$$(X, \ell_X) = \{(0, 0.2), (0, 0.7), (a, 0.2), (a, 0.7), (b, 0.2), (b, 0.7), (c, 0.2), (c, 0.7)\}$$

and it is routine to check that $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ is a two dimensional BCI-eventful algebra. But it is not a two dimensional BCK-eventful algebra since $(0, \ell_0) \otimes (b, \ell_c) = (c, \ell_b) \neq (0, \ell_0)$. Note that $(0, \ell_0) \otimes (0, \ell_b) = (0, \ell_c) = (0, 0.7)$ and $(0, \ell_b) \otimes (0, \ell_0) = (0, \ell_b) = (0, 0.7)$, but $(0, \ell_0) \neq (0, \ell_b)$. Hence $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ does not satisfy the condition (27).

Lemma 3.10. If $X := (X, *, 0)$ is a BCK/BCI-algebra, then $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ is a two dimensional BCK/BCI-eventful algebra.

Proof. Straightforward. □

By a paired BCK/BCI-algebra we mean a two dimensional BCK/BCI-eventful algebra $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ which satisfies the condition (27).

Example 3.11. (1) Consider a BCK-algebra $X = (X, *, 0)$ where $X = \{0, a, b, c\}$ and the binary operation “*” is given by Table 6.

Table 6: Cayley table for the binary operation “*”

*	0	a	b	c
0	0	0	0	0
a	a	0	a	a
b	b	b	0	b
c	c	c	c	0

Given a nonempty set Q , define a mapping ℓ as follows:

$$\ell : X \rightarrow Q, x \mapsto u.$$

Then $(X, \ell_X) = \{(0, u), (a, u), (b, u), (c, u)\}$, and it is clear that $\langle (X, \ell_X), \otimes, (0, u) \rangle$ is a two dimensional BCK-eventful algebra by Lemma 3.10. It is routine to verify that $\langle (X, \ell_X), \otimes, (0, u) \rangle$ satisfies the condition (27). Hence it is a paired BCK-algebra.

(2) Consider the algebra $X = (X, *, 0)$ in Example 3.9 and let ℓ be a mapping from X to a nonempty set Q given by $\ell(x) = v \in Q$ for all $x \in X$. Then $(X, \ell_X) = \{(0, v), (a, v), (b, v), (c, v)\}$, we recall that $X = (X, *, 0)$ is a BCI-algebra, and thus $\langle (X, \ell_X), \otimes, (0, v) \rangle$ is a two dimensional BCI-eventful algebra by Lemma 3.10. It is routine to verify that $\langle (X, \ell_X), \otimes, (0, v) \rangle$ satisfies the condition (27). Hence it is a paired BCI-algebra.

We consider a generalization of Example 3.11.

Theorem 3.12. *Let $X = (X, *, 0)$ be a BCK/BCI-algebra. Given a nonempty set Q , let $\ell : X \rightarrow Q$ be a constant mapping, say $\ell(x) = q$ for all $x \in X$. Then $\langle (X, \ell_X), \otimes, (0, q) \rangle$ is a paired BCK/BCI-algebra.*

Proof. Using Lemma 3.10, we know that $\langle (X, \ell_X), \otimes, (0, q) \rangle$ is a two dimensional BCK/BCI-eventful algebra. Let $x, y, a, b \in X$ be such that $(x, \ell_a) \otimes (y, \ell_b) = (0, q)$ and $(y, \ell_b) \otimes (x, \ell_a) = (0, q)$. Then $(0, q) = (x, \ell_a) \otimes (y, \ell_b) = (x * y, \ell_{a*b}) = (x * y, q)$ and $(0, q) = (y, \ell_b) \otimes (x, \ell_a) = (y * x, \ell_{b*a}) = (y * x, q)$. It follows that $x * y = 0$ and $y * x = 0$. Hence $x = y$, and so $(x, \ell_a) = (x, q) = (y, q) = (y, \ell_b)$. This shows that $\langle (X, \ell_X), \otimes, (0, u) \rangle$ satisfies the condition (27). Therefore it is a paired BCK/BCI-algebra. \square

Theorem 3.12 shows that it can induce many other BCK/BCI-algebras from given a BCK/BCI-algebra. These induced BCK/BCI-algebras are isomorphic each other. Thus a BCK/BCI-algebra induce a unique paired BCK/BCI-algebra up to isomorphism.

Theorem 3.13. *Let $X = (X, *, 0)$ be a BCK/BCI-algebra. Given a nonempty set Q , if a mapping $\ell : X \rightarrow Q$ is one-to-one, then $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ is a paired BCK/BCI-algebra.*

Proof. Using Lemma 3.10, we know that $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ is a two dimensional BCK/BCI-eventful algebra. Let $x, y, a, b \in X$ be such that $(x, \ell_a) \otimes (y, \ell_b) = (0, \ell_0)$ and $(y, \ell_b) \otimes (x, \ell_a) = (0, \ell_0)$. Then $(0, \ell_0) = (x, \ell_a) \otimes (y, \ell_b) = (x * y, \ell_{a*b})$ and $(0, \ell_0) = (y, \ell_b) \otimes (x, \ell_a) = (y * x, \ell_{b*a})$. It follows that $x * y = 0$, $y * x = 0$, $\ell_{a*b} = \ell_0$ and $\ell_{b*a} = \ell_0$. Since ℓ is one-to-one, we have $a * b = 0$ and $b * a = 0$. It follows that $x = y$ and $a = b$. Hence $(x, \ell_a) = (y, \ell_b)$. This shows that $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ satisfies the condition (27). Therefore it is a paired BCK/BCI-algebra. \square

Theorem 3.14. *Let $X = (X, *, 0)$ be a BCK/BCI-algebra. Given a nonempty set Q , if a mapping $\ell : X \rightarrow Q$ satisfies $\ell^{-1}(\ell_0) = \{0\}$, then $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ is a paired BCK/BCI-algebra.*

Proof. Using Lemma 3.10, we know that $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ is a two dimensional BCK/BCI-eventful algebra. Let $x, y, a, b \in X$ be such that $(x, \ell_a) \otimes (y, \ell_b) = (0, \ell_0)$ and $(y, \ell_b) \otimes (x, \ell_a) = (0, \ell_0)$. Then $(0, \ell_0) = (x, \ell_a) \otimes (y, \ell_b) = (x * y, \ell_{a*b})$ and $(0, \ell_0) = (y, \ell_b) \otimes (x, \ell_a) = (y * x, \ell_{b*a})$. It follows that $x * y = 0$, $y * x = 0$, $\ell_{a*b} = \ell_0$ and $\ell_{b*a} = \ell_0$. Hence $a * b \in \ell^{-1}(\ell_0) = \{0\}$ and $b * a \in \ell^{-1}(\ell_0) = \{0\}$ which shows that $a * b = 0$ and $b * a = 0$. It follows that $x = y$ and $a = b$. Hence $(x, \ell_a) = (y, \ell_b)$. This shows that $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ satisfies the condition (27). Therefore it is a paired BCK/BCI-algebra. \square

Lemma 3.15 ([3]). *Given a BCI-algebra $X = (X, *, 0)$, the following are equivalent,*

- (i) X is p -semisimple.
- (ii) $x * (0 * y) = y * (0 * x)$ for all $x, y \in X$.

Theorem 3.16. *The two dimensional BCI-eventful algebra $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ induced by a p -semisimple BCI-algebra $X = (X, *, 0)$ is a commutative group under the operation \odot which is given by*

$$\odot : (X, \ell_X) \times (X, \ell_X) \rightarrow (X, \ell_X), ((x, \ell_a), (y, \ell_b)) \mapsto (x, \ell_a) \otimes ((0, \ell_0) \otimes (y, \ell_b)).$$

Proof. By Lemma 3.10, we know that $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ is a two dimensional BCI-eventful algebra. Let $(x, \ell_a), (y, \ell_b), (z, \ell_c) \in (X, \ell_X)$. Then

$$\begin{aligned} (x, \ell_a) \odot (y, \ell_b) &= (x, \ell_a) \otimes ((0, \ell_0) \otimes (y, \ell_b)) = (x, \ell_a) \otimes (0 * y, \ell_{0*b}) \\ &= (x * (0 * y), \ell_{x*(0*b)}) = (y * (0 * x), \ell_{b*(0*a)}) \\ &= (y, \ell_b) \otimes (0 * x, \ell_{0*a}) = (y, \ell_b) \otimes ((0, \ell_0) \otimes (x, \ell_a)) \\ &= (y, \ell_b) \odot (x, \ell_a) \end{aligned} \tag{28}$$

by Lemma 3.15, and

$$\begin{aligned}
& ((y, \ell_b) \odot (z, \ell_c)) \odot (x, \ell_a) \\
&= ((y, \ell_b) \otimes ((0, \ell_0) \otimes (z, \ell_c))) \otimes ((0, \ell_0) \otimes (x, \ell_a)) \\
&= ((y * (0 * z)) * (0 * x), \ell_{(b*(0*c))*(0*a)}) \\
&= ((y * (0 * x)) * (0 * z), \ell_{(b*(0*a))*(0*c)}) \\
&= ((y, \ell_b) \otimes ((0, \ell_0) \otimes (x, \ell_a))) \otimes ((0, \ell_0) \otimes (z, \ell_c)) \\
&= ((y, \ell_b) \odot (x, \ell_a)) \odot (z, \ell_c).
\end{aligned} \tag{29}$$

Using (28) and (29), we get

$$(x, \ell_a) \odot ((y, \ell_b) \odot (z, \ell_c)) = ((y, \ell_b) \odot (z, \ell_c)) \odot (x, \ell_a) = ((y, \ell_b) \odot (x, \ell_a)) \odot (z, \ell_c) = ((x, \ell_a) \odot (y, \ell_b)) \odot (z, \ell_c).$$

Now,

$$\begin{aligned}
(0, \ell_0) \odot (x, \ell_a) &= (0, \ell_0) \otimes ((0, \ell_0) \otimes (x, \ell_a)) \\
&= (0, \ell_0) \otimes (0 * x, \ell_{0*a}) \\
&= (0 * (0 * x), \ell_{0*(0*a)}) = (x, \ell_a),
\end{aligned}$$

which shows that $(0, \ell_0)$ is the identity element of (X, ℓ_X) . Finally, we show that $(0, \ell_0) \otimes (x, \ell_a)$ is the inverse of any element (x, ℓ_a) . In fact,

$$\begin{aligned}
(x, \ell_a) \odot ((0, \ell_0) \otimes (x, \ell_a)) &= (x, \ell_a) \otimes ((0, \ell_0) \otimes ((0, \ell_0) \otimes (x, \ell_a))) \\
&= (x * (0 * (0 * x)), \ell_{a*(0*(0*a))}) \\
&= (x * x, \ell_{a*a}) = (0, \ell_0).
\end{aligned}$$

Therefore $\langle (X, \ell_X), \odot, (0, \ell_0) \rangle$ is a commutative group. \square

Corollary 3.17. *Let $X = (X, *, 0)$ be a BCI-algebra which satisfies any one of the following assertions.*

$$(\forall x \in X)(0 * x = 0 \Rightarrow x = 0), \tag{30}$$

$$(\forall a \in X)(X = \{a * x \mid x \in X\}), \tag{31}$$

$$(\forall a, x \in X)(a * (a * x) = x), \tag{32}$$

$$(\forall a, x, y, z \in X)((x * y) * (z * a) = (x * z) * (y * a)), \tag{33}$$

$$(\forall x, y \in X)(0 * (y * x) = x * y), \tag{34}$$

$$(\forall x, y, z \in X)((x * y) * (x * z) = z * y). \tag{35}$$

Then the two dimensional BCI-eventful algebra $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ is a commutative group under the operation \odot .

Theorem 3.18. *Let $f : X \rightarrow Y$ be an onto homomorphism of BCK/BCI-algebras. If $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ satisfies the condition (27), then $\langle (Y, \zeta_Y), \otimes, (0, \zeta_0) \rangle$ is a paired BCK/BCI-algebra where ζ is a mapping from Y to Q .*

Proof. By Lemma 3.10, we know that $\langle (X, \ell_X), \otimes, (0, \ell_0) \rangle$ is a two dimensional BCK/BCI-eventful algebra. Let $(x', \zeta_{a'}), (y', \zeta_{b'}), (z', \zeta_{c'}) \in (Y, \zeta_Y)$. Then there exist $x, y, z, a, b, c \in X$ such that $f(x) = x', f(y) = y', f(z) = z', f(a) = a', f(b) = b'$ and $f(c) = c'$. Hence

$$\begin{aligned}
& (((x', \zeta_{a'}) \otimes (y', \zeta_{b'})) \otimes ((x', \zeta_{a'}) \otimes (z', \zeta_{c'}))) \otimes ((z', \zeta_{c'}) \otimes (y', \zeta_{b'})) \\
&= (((f(x), \zeta_{f(a)}) \otimes (f(y), \zeta_{f(b)})) \otimes ((f(x), \zeta_{f(a)}) \otimes (f(z), \zeta_{f(c)}))) \otimes ((f(z), \zeta_{f(c)}) \otimes (f(y), \zeta_{f(b)})) \\
&= ((f(x) * f(y), \zeta_{f(a)*f(b)}) \otimes (f(x) * f(z), \zeta_{f(a)*f(c)})) \otimes (f(z) * f(y), \zeta_{f(c)*f(b)}) \\
&= ((f(x * y), \zeta_{f(a*b)}) \otimes (f(x * z), \zeta_{f(a*c)})) \otimes (f(z * y), \zeta_{f(c*b)}) \\
&= (f(x * y) * f(x * z), \zeta_{f(a*b)*f(a*c)}) \otimes (f(z * y), \zeta_{f(c*b)}) \\
&= (f(((x * y) * (x * z)) * (z * y)), \zeta_{f(((a*b)*(a*c))*(c*b))}) \\
&= (f(0), \zeta_{f(0)}) = (0, \zeta_0),
\end{aligned}$$

$$\begin{aligned}
& (((x', \zeta_{a'}) \otimes (x', \zeta_{a'}) \otimes (y', \zeta_{b'})) \otimes (y', \zeta_{b'})) \\
&= (((f(x), \zeta_{f(a)}) \otimes ((f(x), \zeta_{f(a)}) \otimes (f(y), \zeta_{f(b)}))) \otimes (f(y), \zeta_{f(b)})) \\
&= ((f(x), \zeta_{f(a)}) \otimes (f(x) * f(y), \zeta_{f(a)*f(b)})) \otimes (f(x), \zeta_{f(a)}) \\
&= ((f(x), \zeta_{f(a)}) \otimes (f(x * y), \zeta_{f(a*b)})) \otimes (f(x), \zeta_{f(a)}) \\
&= (f(x * (x * y)), \zeta_{f(a*(a*b))}) \otimes (f(x), \zeta_{f(a)}) \\
&= (f((x * (x * y)) * y), \zeta_{f((a*(a*b))*b)}) \\
&= (f(0), \zeta_{f(0)}) = (0, \zeta_0),
\end{aligned}$$

and $(x', \zeta_{a'}) \otimes (x', \zeta_{a'}) = (f(x), \zeta_{f(a)}) \otimes (f(x), \zeta_{f(a)}) = (f(x * x), \zeta_{f(a*a)}) = (f(0), \zeta_{f(0)}) = (0, \zeta_0)$. Hence $\langle (Y, \zeta_Y), \otimes, (0, \zeta_0) \rangle$ is a two dimensional BCI-eventful algebra. Since $(0, \zeta_0) \otimes (x', \zeta_{a'}) = (f(0), \zeta_{f(0)}) \otimes (f(x), \zeta_{f(a)}) = (f(0 * x), \zeta_{f(0*a)}) = (f(0), \zeta_{f(0)}) = (0, \zeta_0)$, we know that $\langle (Y, \zeta_Y), \otimes, (0, \zeta_0) \rangle$ is a two dimensional BCK-eventful algebra. Assume that $(x', \zeta_{a'}) \otimes (y', \zeta_{b'}) = (0, \zeta_0)$ and $(y', \zeta_{b'}) \otimes (x', \zeta_{a'}) = (0, \zeta_0)$. Then

$$(0, \zeta_0) = (x', \zeta_{a'}) \otimes (y', \zeta_{b'}) = (f(x), \zeta_{f(a)}) \otimes (f(y), \zeta_{f(b)}) = (f(x) * f(y), \zeta_{f(a)*f(b)})$$

and

$$(0, \zeta_0) = (y', \zeta_{b'}) \otimes (x', \zeta_{a'}) = (f(y), \zeta_{f(b)}) \otimes (f(x), \zeta_{f(a)}) = (f(y) * f(x), \zeta_{f(b)*f(a)}),$$

which imply that $f(x) * f(y) = 0$, $f(y) * f(x) = 0$, $f(a) * f(b) = 0$ and $f(b) * f(a) = 0$. Hence $x' = f(x) = f(y) = y'$ and $a' = f(a) = f(b) = b'$. Therefore $(x', \zeta_{a'}) = (y', \zeta_{b'})$. Consequently, $\langle (Y, \zeta_Y), \otimes, (0, \zeta_0) \rangle$ is a paired BCK/BCI-algebra. \square

4 Conclusions

We have introduced two-dimensional event sets and have applied it to algebraic structures. We have introduced the notions of two dimensional BCK/BCI-eventful algebra, paired B-algebra and paired BCK/BCI-algebra, and have investigated several properties. We have considered conditions for two dimensional eventful algebra to be a B-algebra and a BCK/BCI-algebra. We have discussed the process of inducing a paired B-algebra using a group, and have established a commutative group using two dimensional BCI-eventful algebra. We have presented examples to show that a two dimensional eventful BCK/BCI-algebra is not a BCK/BCI-algebra, and then we have considered conditions for a two dimensional eventful BCK/BCI-algebra to be a BCK/BCI-algebra. We have studied a paired BCK/BCI-algebra in relation to the BCK/BCI-homomorphism.

References

- [1] A.A.A. Agboola, B. Davvaz, F. Smarandache, *Neutrosophic quadruple algebraic hyperstructures*, Annals of Fuzzy Mathematics and Informatics, 14(1) (2017), 29–42.
- [2] S.A. Akinleye, F. Smarandache, A.A.A. Agboola, *On neutrosophic quadruple algebraic structures*, Neutrosophic Sets and Systems, 12 (2016), 122–126.
- [3] Y. Huang, *BCI-algebra*, Science Press, Beijing, 2006.
- [4] Y.B. Jun, S.Z. Song, S.J. Kim, *Event and its application in algebraic structures*, New Mathematics and Natural Computation, (in press).
- [5] Y.B. Jun, S.Z. Song, F. Smarandache, H. Bordbar, *Neutrosophic quadruple BCK/BCI-algebras*, Axioms, 7 (2018), 41.
- [6] J. Meng, Y.B. Jun, *BCK-algebras*, Kyungmoonsa Co. Seoul, Korea, 1994.

- [7] M. Mohseni Takallo, Y.B. Jun, *Commutative neutrosophic quadruple ideals of neutrosophic quadruple BCK-algebras*, Journal of Algebraic Hyperstructures and Logical Algebras, 1(1) (2020), 95–105.
- [8] J. Neggers, H.S. Kim, *On B-algebras*, Matematicki Vesnik, 54(1-2) (2002), 21–29.
- [9] F. Smarandache, *Neutrosophy, neutrosophic probability, set, and logic*, Proquest Information and Learning, Ann Arbor, Michigan, USA, 105 p., 1998. <http://fs.gallup.unm.edu/eBook-neutrosophics6.pdf> (last edition online).
- [10] F. Smarandache, *A unifying field in logics: Neutrosophic logic. Neutrosophy, neutrosophic set, neutrosophic probability*, American Reserch Press, Rehoboth, NM, 1999.
- [11] F. Smarandache, *Neutrosophic quadruple numbers, refined neutrosophic quadruple numbers, absorbance law, and the multiplication of neutrosophic quadruple numbers*, Neutrosophic Sets and Systems, 10 (2015), 96–98.